

ORIGINAL ARTICLE

Open Access

# Insights into bacterial CO<sub>2</sub> metabolism revealed by the characterization of four carbonic anhydrases in *Ralstonia eutropha* H16

Claudia S Gai<sup>1</sup>, Jingnan Lu<sup>2</sup>, Christopher J Brigham<sup>1,3</sup>, Amanda C Bernardi<sup>1</sup> and Anthony J Sinskey<sup>1,4,5\*</sup>

## Abstract

Carbonic anhydrase (CA) enzymes catalyze the interconversion of CO<sub>2</sub> and bicarbonate. These enzymes play important roles in cellular metabolism, CO<sub>2</sub> transport, ion transport, and internal pH regulation. Understanding the metabolic role of CAs in the chemolithoautotrophic bacterium *Ralstonia eutropha* is important for the development of high performance fermentation processes based on the bacterium's capability to fix carbon using the Calvin-Benson-Bassham (CBB) cycle. Analysis of the *R. eutropha* H16 genome sequence revealed the presence of four CA genes: *can*, *can2*, *caa* and *cag*. We evaluated the importance of each of the CAs in the metabolism of *R. eutropha* by examination of growth and enzyme activity in gene deletion, complementation, and overexpression strains. All four purified CAs were capable of performing the interconversion of CO<sub>2</sub> and HCO<sub>3</sub><sup>-</sup>, although the equilibrium towards the formation of CO<sub>2</sub> or HCO<sub>3</sub><sup>-</sup> differs with each CA. Deletion of *can*, encoding a β-CA, affected the growth of *R. eutropha*; however the growth defect could be compensated by adding CO<sub>2</sub> to the culture. Deletion of the *caa*, encoding an α-CA, had the strongest deleterious influence on cell growth. Strains with deletion or overexpression of *can2* or *cag* genes exhibited similar behavior to wild type under most of the conditions tested. In this work, Caa was studied in greater detail using microscopy and complementation experiments, which helped confirm its periplasmic localization and determine its importance for robust growth of *R. eutropha*. A hypothesis for the coordinated role of these four enzymes in the metabolism of *R. eutropha* is proposed.

**Keywords:** CO<sub>2</sub> transport; *Cupriavidus necator*; Carbon dioxide; Bicarbonate; Dissolved inorganic carbon; Periplasm; Zinc metalloenzyme

## Introduction

Carbon dioxide, bicarbonate, carbonic acid, and carbonate are key metabolites in all living systems, and the equilibrium of these different forms in living cells is important for proper physiological functioning. Carbonic anhydrase (CA) (EC 4.2.1.1) catalyzes the interconversion between carbon dioxide and bicarbonate (CO<sub>2</sub> + H<sub>2</sub>O ↔ HCO<sub>3</sub><sup>-</sup> + H<sup>+</sup>). The CA-catalyzed metabolic hydration/dehydration of CO<sub>2</sub> and HCO<sub>3</sub><sup>-</sup> supports physiological functions in the cell, even though the uncatalyzed (enzyme-free) reaction proceeds at significant rates [1].

There are five main classes of CA enzyme found in nature (α, β, γ, δ and ζ). There are no significant sequence similarities observed between representatives of the different classes, implying that CA enzymes evolved convergently in diverse biological systems [2,3]. The α-CA enzymes are the most studied to date and are found in many different genera. They are monomeric, zinc-containing enzymes with multiple isozymes, which have different expression patterns and localizations [3-6]. In α-CA enzymes, Zn<sup>2+</sup> is coordinated by three amino acids (usually histidines) and a water molecule. The role of Zn<sup>2+</sup> is to facilitate the deprotonation of water for the formation of a nucleophilic hydroxide ion. This hydroxide ion can then initiate a nucleophilic attack on the carbonyl group of CO<sub>2</sub> and convert it into bicarbonate. Specifically, the +2 charge of the zinc ion associates with the oxygen atom of water, supplying positive charge to

\* Correspondence: asinskey@mit.edu

<sup>1</sup>Department of Biology, Massachusetts Institute of Technology, Bldg. 68-370, Cambridge, MA 02139, USA

<sup>4</sup>Division of Health Sciences and Technology, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Bldg. 68-370, Cambridge, MA 02139, USA

Full list of author information is available at the end of the article

the hydroxide ion, which is then able to attack the CO<sub>2</sub> [2]. All CA classes present differences in their secondary, tertiary and quaternary structures. Enzymes of the  $\alpha$  class are monomeric,  $\beta$ -CA enzymes are usually oligomeric and contain 2 to 6 monomers, and  $\gamma$ -CA enzymes associate as homotrimers. Also, the active site ligands are potentially different among the CA classes, although  $\beta$ -CA and  $\gamma$ -CA enzymes are believed to function in a manner similar to  $\alpha$ -CAs, in which both use the same zinc hydroxide mechanism [7].

*Ralstonia eutropha* (also *Cupriavidus necator*) is a Gram-negative facultative chemoautotrophic betaproteobacterium. It is well known for its ability to produce polyhydroxyalkanoate under high carbon but limited nitrogen or phosphorus conditions [8,9]. The capacity of *R. eutropha* to grow autotrophically using CO<sub>2</sub> as the sole carbon source has been recently explored and studied for the production of alternative biofuels [10,11]. Assimilation of CO<sub>2</sub> during autotrophic growth of *R. eutropha* proceeds by the Calvin-Benson-Bassham (CBB) cycle [12] and requires large amounts of energy to fuel the synthesis of cellular building blocks. Organisms must have a reliable and efficient system of controlling intracellular pH and CO<sub>2</sub> concentrations in order to carry out carbon fixation [13]. Cyanobacteria evolved carboxysomes as an efficient mechanism to increase CO<sub>2</sub> concentration and consequently its fixation efficiency [14,15], but *R. eutropha* lacks this system although it contains its two main enzymes, CA and RuBisCO [16]. Besides the key CBB cycle enzyme, RuBisCO, CA is of great importance for fine-tuning the concentration of CO<sub>2</sub> in autotrophic metabolism.

Four putative CA genes were identified in the genome sequence of *R. eutropha* strain H16. Two CA genes are located on chromosome 1, and the others are on chromosome 2. The *can* (locus tag H16\_A0169) and *can2* (locus tag H16\_B2270) genes encode  $\beta$ -CA enzymes, the *caa* (locus tag H16\_B2403) gene encodes a putative periplasmic  $\alpha$ -CA, and the gene with locus tag H16\_A1192 (hereafter known as *cag*) encodes a  $\gamma$ -like-CA/acetyltransferase [16]. The presence of CA genes of multiple classes in *R. eutropha* suggests that the gene products play major roles in CO<sub>2</sub> transport and metabolism. Additionally, the diversity of CA gene products expressed in *R. eutropha* implies that the functions of these different enzymes could all be unique. Dobrinski et al. [17] examined four CA enzymes ( $\alpha$ ,  $\beta$ ,  $\gamma$  and CsoSCA) from the deep sea proteobacterium *Thiomicrospira crunogena* and suggested different roles for each of the enzymes in relation to carbon fixation capabilities and survival mechanisms of the microorganism. Currently, the exact roles of all four *R. eutropha* CA enzymes are still largely

unknown and the only one studied in depth to date is Can, which was identified as being essential for growth under atmospheric concentrations of CO<sub>2</sub> [1].

In the present study, we examined the activities of all four CA enzymes from *R. eutropha*, following heterologous expression and purification from *Escherichia coli*. We also assessed the effects of single and combinatorial CA gene deletions on cell physiology and fitness. The importance of Caa localization in the cell was further examined by overexpressing the enzyme with and without a periplasmic localization signal peptide in a  $\Delta caa$  strain. Periplasmic localization was confirmed by detection of a fusion protein of Red Fluorescent Protein (RFP) and Caa using fluorescent microscopy.

## Materials and methods

### Chemicals, bacterial strains and plasmids

Chemicals were purchased from Sigma-Aldrich unless indicated otherwise. Experiments were performed with strains and plasmids listed in Table 1.

### Growth media and cultivation conditions

*R. eutropha* strains were propagated in tryptic soy broth (TSB) (Becton Dickinson, Sparks, MD) or minimal medium [11] with fructose at a final concentration of 1% or 2% (w vol<sup>-1</sup>), or pyruvate, lactate, succinate, or formate, each at a final concentration of 0.2% (w vol<sup>-1</sup>) or (vol vol<sup>-1</sup>). All cultures were inoculated to an initial OD<sub>600nm</sub> of 0.05. *E. coli* strains were grown in LB medium [21] at 37°C. For growth experiments in a CO<sub>2</sub>-rich environment, cultures were performed inside a CO<sub>2</sub> incubator (Napco 6100 - Thermo Electron Corporation, Winchester, VA USA) with an atmosphere of 10% CO<sub>2</sub> at 30°C, under 200 rpm agitation. Appropriate antibiotics were added to the growth media at the following concentrations: gentamicin, 10  $\mu$ g mL<sup>-1</sup>; kanamycin, 200  $\mu$ g mL<sup>-1</sup> (for *R. eutropha*); kanamycin, 50  $\mu$ g mL<sup>-1</sup> (for *E. coli*); ampicillin, 100  $\mu$ g mL<sup>-1</sup>; chloramphenicol, 34  $\mu$ g mL<sup>-1</sup>.

Autotrophic cultures were prepared using minimal media without carbon source. For these cultures, a 250-mL Erlenmeyer flask containing 30 mL of culture is placed into a 15000 Vacu-Quik Jar System (Almore International, Inc., Portland, OR, USA). Agitation inside the flask was assured by a stir bar. Vacuum and nitrogen were applied alternatively and repeatedly three times to flush the chamber and remove all air. An approximate molar ratio of 8:1:1 (H<sub>2</sub>: O<sub>2</sub>: CO<sub>2</sub>) [22] gas mixture was then supplied to the jar. Any additional space in the jar was filled with 100% nitrogen.

**Table 1 Plasmids and bacterial strains used in this work with relevant genotype characteristics**

Strain or plasmid	Relevant characteristics	Reference or source
<i>Ralstonia eutropha</i>		
H16	Wild-type gentamycin resistant (Gen <sup>r</sup> )	ATCC 17699
Re2061	H16 $\Delta$ <i>phaCAB</i> (Gen <sup>r</sup> )	[11]
Re2427	H16 $\Delta$ <i>can</i>	This study
Re2428	H16 $\Delta$ <i>caa</i>	This study
Re2430	H16 $\Delta$ <i>cag</i>	This study
Re2437	H16 $\Delta$ <i>can2</i>	This study
Re2436	H16 $\Delta$ <i>can</i> $\Delta$ <i>can2</i> $\Delta$ <i>caa</i> $\Delta$ <i>cag</i>	This study
<b><i>Escherichia coli</i></b>		
DH10-beta competent cells	Strain suitable for high efficiency transformation	New England Biolabs
BL21(DE3)	Strain suitable for transformation and protein expression	New England Biolabs
S17-1	Conjugation strain for transfer of plasmids into <i>R. eutropha</i>	[18]
<b>Plasmids</b>		
pBBR1MCS-2	Broad-host-range cloning vector confers kanamycin resistance (Kan <sup>r</sup> )	[19]
pCan	pBBR1MCS-2 containing <i>can</i> gene (H16 A0169) (Kan <sup>r</sup> )	This work
pCan2	pBBR1MCS-2 containing <i>can2</i> gene (H16 B2270) (Kan <sup>r</sup> )	This work
pCaa	pBBR1MCS-2 containing <i>caa</i> gene (H16 B2403) (Kan <sup>r</sup> )	This work
pCaaB	pBBR1MCS-2 containing <i>caa</i> gene without the <i>N</i> terminal signaling peptide sequence (H16 B2403) (Kan <sup>r</sup> )	This work
pCag	pBBR1MCS-2 containing <i>cag</i> gene (H16 A1192) (Kan <sup>r</sup> )	This work
pETCan	pET14b containing <i>can</i> gene (H16 A0169) (Amp <sup>r</sup> )	This work
pETCan2	pET14b containing <i>can2</i> gene (H16 B2270) (Amp <sup>r</sup> )	This work
pETCaa	pET14b containing <i>caa</i> gene (H16 B2403) (Amp <sup>r</sup> )	This work
pETCag	pET14b containing <i>cag</i> gene (H16 A1192) (Amp <sup>r</sup> )	This work
pStrepCan	pET51b containing <i>can</i> gene (H16 A0169) (Amp <sup>r</sup> )	This work
pStrepCaa	pET51b containing <i>caa</i> gene (H16 B2403) (Amp <sup>r</sup> )	This work
pStrepCaaB	pET51b containing <i>caa</i> gene without the <i>N</i> terminal signaling peptide sequence (H16 B2403) (Amp <sup>r</sup> )	This work
pRARE	Overcoming the codon bias of <i>E. coli</i> for enhanced protein expression (Cam <sup>r</sup> )	[20]
pJV7	pJQ200Kan with <i>phaC1</i> inserted into <i>Bam</i> HI restriction site, confers kanamycin resistance (Kan <sup>r</sup> )	[9]
pJV7 $\Delta$ <i>can</i>	pJV7 with <i>phaC1</i> allele removed by <i>Bam</i> HI digestion and replaced with $\Delta$ <i>can</i> allele (Kan <sup>r</sup> )	This work
pJV7 $\Delta$ <i>can2</i>	pJV7 with <i>phaC1</i> allele removed by <i>Bam</i> HI digestion and replaced with $\Delta$ <i>can2</i> allele (Kan <sup>r</sup> )	This work
pJV7 $\Delta$ <i>caa</i>	pJV7 with <i>phaC1</i> allele removed by <i>Bam</i> HI digestion and replaced with $\Delta$ <i>caa</i> allele (Kan <sup>r</sup> )	This work
pJV7 $\Delta$ <i>cag</i>	pJV7 with <i>phaC1</i> allele removed by <i>Bam</i> HI digestion and replaced with $\Delta$ <i>cag</i> allele (Kan <sup>r</sup> )	This work
pRFP	pBBR1MCS-2 containing <i>rfp</i> gene amplified from JBp000066 kindly offered by J. Mueller (JBEI) (Kan <sup>r</sup> )	This work
pCAA_RFP	pBBR1MCS-2 containing <i>caa</i> gene fused by a 6aa linker to the <i>rfp</i> gene (Kan <sup>r</sup> )	This work

### Plasmid and strain construction

Standard protocols were employed for DNA manipulation [23]. PCR amplification of DNA was performed using Phusion DNA Polymerase (New England Biolabs, Ipswich, MA, USA). Restriction enzymes and T4 DNA ligase used in this study were from New England Biolabs

(Ipswich, MA, USA). The QIAquick Gel Extraction Kit (Qiagen, Valencia, CA, USA) was used for all gel purifications of DNA products and plasmid extractions were carried out using QIAprep Spin Miniprep Kit.

For the construction of overexpression and complementation plasmids, each of the four CA genes was cloned into

pBBR1MCS-2 between restriction sites *KpnI* and *HindIII* (*can*, *can2* and *caa*) or *Sall* and *HindIII* (*cag*) to create the plasmids pCan, pCan2, pCaa and pCag respectively (Table 1). Each plasmid was transformed by electroporation into *E. coli* S17-1, which was then used as a donor strain for the conjugative transfer of plasmid into *R. eutropha* by a standard mating procedure [24]. For control experiments, *R. eutropha* was also transformed with the empty vector pBBR1MCS-2.

For the construction of plasmids for heterologous overexpression of CA in *E. coli*, followed by protein purification, CA genes were amplified and cloned as described above using the restriction sites with *NdeI* and *XhoI* (*can*, *can2* and *caa*) or *NdeI* and *BamHI* (*cag*) on both plasmids pET14b (*N*-terminal His-tag) and pET51b (*C*-terminal Strep2-tag) (Table 1). Each plasmid was then transformed into *E. coli* BL21 (DE3) harboring pRARE (Table 1).

For the assessment of the role of the predicted Caa signaling peptide and the effect of the correct localization of the enzyme on *R. eutropha* physiology, the entire annotated gene (*caa* -H16\_B2403), with (*caa*) and without (*caaB*) the nucleotide sequence encoding the 23-aa *N*-terminal predicted signal-peptide were amplified and separately cloned into expression vectors (Table 1). The 23-aa *N*-terminal periplasmic signal peptide of Caa was predicted using SignalP 4.0 software [25].

RFP fusion plasmids were prepared by amplifying and inserting the *rfp* gene from the plasmid JBp000066 (kindly provided by J. Mueller and S. Singer – JBEI, Emeryville, CA, USA) at the 3' end of the *caa* gene, with DNA sequence encoding a 6-glycine linker between the genes, into pBBR1MCS-2 using the Gibson assembly method [26] to produce pCaa\_RFP. The *rfp* gene alone was expressed separately as a control (pRFP). Plasmids were introduced separately, as described above, into *R. eutropha* strain Re2061 (Table 1), which does not produce intracellular polyhydroxyalkanoate.

Plasmids for markerless deletion of CA genes were constructed according to Lu et al. [11]. Once the deletion strains Re2427 (H16  $\Delta$ *can*), Re2428 (H16  $\Delta$ *caa*), Re2430 (H16  $\Delta$ *cag*), and Re 2437(H16  $\Delta$ *can2*) were constructed, overexpression plasmids containing *can*, *can2*, *caa* or *cag* genes were introduced *in trans* to create Re2427/pCan, Re2428/pCaa, Re2430/pCag, Re2437/pCan2 and Re2428/pCaaB for complementation studies. A strain with quadruple CA gene deletions, Re2436 (H16 $\Delta$ *can* $\Delta$ *can2* $\Delta$ *caa* $\Delta$ *cag*), was also constructed, and each CA enzyme, expressed *in trans*, was complemented back into the strain to create Re2436/pCan, Re2436/pCaa, Re2436/pCag and Re2436/pCan2. All oligonucleotide primers used for plasmid and strain constructions are listed in the Additional file 1: Table S1.

### Cell preparation and protein purification

*R. eutropha* CA genes were expressed in *E. coli* BL21 (DE3)/pRARE using plasmids listed in Table 1. Cells were induced with IPTG (Isopropyl  $\beta$ -D-1-thiogalactopyranoside, 0.6 mM final concentration) at a culture OD<sub>600</sub> of 0.4-0.6. Cells were harvested 8 h post-induction by centrifugation and stored at -80°C prior to CA purification and activity assay. Purification of Strep2-tagged proteins was performed using a gravity flow Strep-Tactin® Superflow® high capacity column (IBA, Göttingen, Germany) following the manufacturer's instructions. His-tagged proteins were purified using a 5-mL HisTrap® FF column (Amersham Bioscience, Uppsala, Sweden) operated by a low pressure liquid chromatography system (BioRad, Hercules, California, USA). Fractions containing purified CA were detected on a 12% SDS-PAGE gel [23], concentrated using Amicon Ultra-15 Centrifugal Filter Units (EMD Millipore Corporation, Billerica, MA, USA), and dialyzed overnight using a Slide-A-Lyzer Dialysis Cassette (Thermo Scientific, Asheville, NC, USA) in Tris-HCl (100 mM pH 7) at 4°C.

### Determination of CA activity

The CA activity assays were based on methods from Sundaram et al. [27] and Fasseas et al. [4] with some modifications. CA activity was measured by comparing the rapid change of pH in the presence of CA to the slower pH change in the absence of CA. The assays were performed using a stopped-flow device (Applied Photophysics Ltd, Leatherhead, United Kingdom) connected to a spectrophotometer (Agilent 8453 UV-Visible Kinetic Mode). Purified CA enzymes were diluted in assay buffer (50 mM Na<sub>2</sub>SO<sub>4</sub>, 50 mM HEPES, 50 mM MgSO<sub>4</sub>, 0.004% (w v<sup>-1</sup>) Phenol red, pH 8 for assays using CO<sub>2</sub> as a substrate or pH 6 for assays using KHCO<sub>3</sub> as a substrate. All reagents were kept on ice while the spectrophotometer and stopped-flow apparatus were kept at 1°C  $\pm$  1°C using a recirculating water bath. The substrates used for the activity assays were KHCO<sub>3</sub> (50 mM) and saturated CO<sub>2</sub> (dry ice in assay buffer for 30 min). The final volume of the assay reaction was 400  $\mu$ L and the absorbance A<sub>557nm</sub> was measured for 60 sec. Control assays were carried out in the absence of enzymes.

Enzyme Units (EU) were calculated as illustrated below by comparing the time of pH change in the presence of enzyme to reaction without enzyme [27]. Each additional EU speeds the catalytic activity of the enzyme by two fold. The assay was validated using commercially available purified CA from bovine erythrocytes (Sigma-Aldrich C9207). Protein concentration (purified or cell extract) was determined by standard Bradford assay [28].

$$EU = \frac{\text{Time of uncatalyzed} - \text{Time of catalyzed reaction (s)}}{\text{Time of catalyzed reaction (s)}} \times \frac{1}{\text{Total protein in assay (mg)}}$$

Adapted from [27].

### Microscopy

Slides of Re2061/pCaa\_RFP or Re2061/pRFP were prepared using 10  $\mu$ L of overnight TSB cultures placed onto Poly-L-Lysine-coated slides. Cells were observed under 100 $\times$  magnification using a Nikon Labophot-2 microscope with phase-contrast and fluorescence attachments. Images were acquired with a SPOT cooled color digital camera (Diagnostic Instruments, Inc.) and prepared using Adobe Photoshop version 5.0.

### Polymer quantification

Polyhydroxybutyrate (PHB) content was determined as described previously [29,30].

## Results


### Characterization of CA enzymes

All four carbonic anhydrase enzymes were heterologously overexpressed in recombinant *E. coli*, purified, and examined in this work. Each CA was able to catalyze the interconversion reaction between CO<sub>2</sub> and bicarbonate. Additionally, each CA exhibited different specific activity and substrate preference towards CO<sub>2</sub> and bicarbonate. Caa presented a higher activity using CO<sub>2</sub>

as the substrate, compared to the activity of Can, Can2 or Cag, which all preferred bicarbonate (Figure 1). The specific activity of Caa towards CO<sub>2</sub> as a substrate ( $422.2 \pm 97.1$  EU mg<sup>-1</sup>) was higher than the activity measured for the other CAs (Can  $59.5 \pm 15.3$  EU mg<sup>-1</sup>; Can2  $157.9 \pm 4$  EU mg<sup>-1</sup> and Cag  $138.2 \pm 61.3$  EU mg<sup>-1</sup>). On the other hand, the specific activities of both Can and Can2 were similar with bicarbonate as a substrate ( $277.9 \pm 12.6$  EU mg<sup>-1</sup> and  $309.4 \pm 103.1$  EU mg<sup>-1</sup> respectively). Cag presented the highest activity using bicarbonate as a carbon source ( $488.5 \pm 122.1$  EU mg<sup>-1</sup>). All CAs were active towards CO<sub>2</sub> and bicarbonate, however only Caa had a distinct preference for CO<sub>2</sub> as a substrate while the other three CAs all preferred bicarbonate. The tests for validation of the assay were done using a commercial CA and the values obtained were  $130.7 \pm 20.1$  EU mg<sup>-1</sup> using bicarbonate as a substrate and  $168.8 \pm 5.9$  EU mg<sup>-1</sup> using CO<sub>2</sub> as a substrate. Comparative CO<sub>2</sub> hydration values reported for CAs of different microorganisms can be found on Additional file 1: Table S2.

### Growth characterization of overexpression strains

The growth of CA overexpression strains H16/pCan, H16/pCan2, H16/pCaa and H16/pCag were tested in



**Figure 1 Specific activity of purified Carbonic Anhydrases.** Specific activity of purified, His-tagged carbonic anhydrase enzymes. Substrates used for this assay were bicarbonate (50 mM, dark bars) and CO<sub>2</sub> (saturated solution, grey bars). CA activity was measured by comparing the rapid change of pH in the presence of CA to the slower pH change in the absence of CA. The pH change was monitored using phenol red and the absorbance  $A_{557nm}$  was measured for 60 sec. Average values from three experiments were plotted with error bars representing the standard deviation.




minimal medium with 2% (w vol<sup>-1</sup>) fructose in the presence of ambient CO<sub>2</sub> concentrations (air) and under autotrophic condition (H<sub>2</sub>:CO<sub>2</sub>:O<sub>2</sub>). Using fructose as the carbon source, all strains exhibited similar growth behavior (Figure 2A)

CA activity of cell extracts of *R. eutropha* strains grown on fructose was determined. As shown in Figure 2B, the CA-overexpression strains exhibited higher activity than the control using both substrates. Notably high CA activity was observed in H16/pCaa using CO<sub>2</sub> as a substrate. Using bicarbonate as a substrate, all the strains presented significantly higher activity than the control (P < 0.01). However, each CA overexpression strain exhibited no significant difference in activity using bicarbonate as the substrate. These

results indicate the preference of Caa for CO<sub>2</sub> as a substrate, which is in accordance with activity results using purified Caa (Figure 1). The PHB content of the CA overexpression and control strains was also assessed, and it was observed that all strains grown using fructose as the main carbon source produced similar intracellular amounts of PHB (data not shown).

Under autotrophic conditions, significant differences in growth were observed between the different strains. Growth of H16/pCaa was significantly lower than all the other strains (P < 0.01). H16/pCag grew to a greater extent compared to the others (P < 0.05), while H16/pCan and H16/pCan2 presented no significant growth difference compared to H16/pBBR1MCS-2 (Figure 2C).



**Figure 2 Growth, CA activity and PHB production of overexpression strains. A)** Growth of CA overexpression strains H16/pCan, H16/pCan2, H16/pCaa and H16/pCag, compared to the control strain (H16/pBBR1MCS-2) with 2% (w vol<sup>-1</sup>) fructose as the main carbon source. **B)** Carbonic anhydrase activity of overexpression strains after 24 h of growth in minimal medium with 2% (w vol<sup>-1</sup>) fructose. H16/pBBR1MCS-2 was used as a control. The Tukey test was used as statistic test to compare all the strains (\*) represents significant difference in activity using CO<sub>2</sub> as a substrate. (\*\*) represents significant difference in activity using bicarbonate, P < 0.01. **C)** Growth of *R. eutropha* strains H16/pCan, H16/pCan2, H16/pCaa and H16/pCag and control strain H16/pBBR1MCS-2 under autotrophic conditions. (\*) represents significant difference in final OD<sub>600nm</sub> P < 0.01. (\*\*) represents significant difference in final OD<sub>600nm</sub>, P < 0.05. **D)** PHB production in 48 h by the overexpression and control strains and under autotrophic conditions. (\*) represents significant difference in PHB production per cell dry weight (CDW) (w w<sup>-1</sup>), P < 0.01. Values represented were averages from three replicates with standard deviation represented by error bars.

These strains were tested for PHB production after 48 h of autotrophic growth (Figure 2D). H16/pCaa produced the lowest amount of PHB per CDW, compared to the other CA overexpression and control strains.

### Growth characteristics of the CA deletion strains


The deletion strains (Re2427, Re2428, Re2430 and Re2437) were cultivated in minimal medium supplemented with various carbon sources (fructose, formate, pyruvate, succinate, or lactate) in the presence of air or air supplemented with 10% CO<sub>2</sub>. Strain Re2428 (H16Δcaa) was unable to grow under all conditions tested, except in fructose with excess CO<sub>2</sub>, where it grew slightly to a final OD<sub>600nm</sub> of 0.2 (Figure 3F). Growth of strain Re2427 (H16Δcan) was completely recovered with the addition of CO<sub>2</sub> to the fructose culture (Figure 3F), although this strain grew poorly under other nutrient-supplemented conditions (Figure 3A-E). Strain Re2437 (H16Δcan2) grew no differently than the wild type, except in the presence of fructose supplemented with CO<sub>2</sub>, in which the growth output dropped by nearly 60%. Re2437 cells presented different morphology under light microscopy without CO<sub>2</sub> supplement. These cells were longer than the wild type cells under such condition, (Additional file 1: Figure S2) which can be a sign of stress that was not detected during absorbance measurements and could be caused by a pH imbalance during growth. Additional file 1: Figure S2 shows light

microscopy images of the deletion strains. In the presence of increased CO<sub>2</sub> concentrations, increasing stress could be detected by an observed decrease in absorbance (Figure 3F). This suggests the importance of Can2 in maintaining the cellular pH, because the addition of CO<sub>2</sub> to the media decreases the pH in the media, affecting consequently the cytoplasmic pH. To test this, strain Re2437 was cultivated in media under different pH values (5.5, 7.0 and 8.5). The growth of Re2437 was less affected by this pH change as compared to wild type. Growth of *R. eutropha* H16 and Re2427 with different initial pH values can be seen in Additional file 1: Figure S3. The growth of Re2430 (H16Δcag) was similar to wild type under all conditions tested.

### Investigation of the phenotypic effects of individual CA enzymes

The quadruple mutant, Re2436 (H16 Δcan Δcan2 Δcaa Δcag), had each of the four CA-encoding genes inserted *in trans* to create Re2436/pCan, Re2436/pCaa, Re2436/pCag, and Re2436/pCan2 (see Materials and methods), in order to demonstrate the isolated effect of each CA on *R. eutropha* metabolism.

As expected, none of the strains completely recovered the phenotype when growing on minimal media with fructose as a carbon source in air or CO<sub>2</sub>-enriched environment (data not shown). In ambient air condition, no significant difference was observed with growth of all



four strains compared to wild type (data not shown). On the other hand, with the addition of 10% CO<sub>2</sub>, Re2436/pCan exhibited a higher growth rate when compared to Re2436 and the other strains (Figure 4).

The CA single deletion strains were transformed with the respective plasmid to complement the CA deletion to obtain Re2427/pCan, Re2428/pCaa, Re2430/pCag and Re2437/pCan2. When tested on minimal media with fructose 2% (w vol<sup>-1</sup>) the strains Re2427/pCan, Re2430/pCag and Re2437/pCan2 exhibited growth comparable to wild type. Growth of complemented deletion strains can be seen in Additional file 1: Figure S1. The only exception was Re2428/pCaa, which was unable to grow to an extent comparable with the wild type under any condition tested and was therefore the subject of closer examination in this study.


### A closer look at Caa

Examination of the  $\alpha$ -periplasmic enzyme, Caa, yielded very interesting results during this study. Besides being the only  $\alpha$ -carbonic anhydrase identified in the genome sequence of *R. eutropha* strain H16, it is described as a “putative periplasmic enzyme.” In this study, the purified Caa is the only *R. eutropha* CA capable of performing the interconversion of CO<sub>2</sub> and HCO<sub>3</sub><sup>-</sup> with equilibrium lying towards the formation of HCO<sub>3</sub><sup>-</sup> (Figure 1). The results of Caa overexpression in *R. eutropha* H16 grown in autotrophic culture indicated that the overexpression of Caa in the wild type strain had a negative influence on growth (Figure 2C) and PHB production (Figure 2D). Moreover, in any medium tested, the *caa* deletion strain,

Re2428, was unable to recover growth to wild type levels (Figure 3).


In order to study the importance of Caa cellular localization on the phenotype recovery of Re2428, *caa* was cloned both as annotated and without the *N*-terminal signaling peptide, which hypothetically would prevent it from being targeted to the periplasm of *R. eutropha*. Purified, Strep2-tagged Caa enzyme without the signaling peptide (CaaB) exhibited similar activity values when compared to the purified, Strep-tagged Caa enzyme containing the signaling peptide. Purified Can was used as a control for CA activity assay (Figure 5A).

The *caa* and *caaB* genes were reintroduced *in trans* into Re2428 using plasmids pCaa and pCaaB (Table 1), in an attempt to recover *R. eutropha* growth in air and to determine the effect of Caa localization on cell growth. *R. eutropha* wild type, Re2428, and complemented strains (Re2428/pCaa and Re2428/pCaaB) were cultivated in TSB and minimal media. The results presented in Figure 5B and C show that Re2428/pCaa and Re2428/pCaaB grew poorly under the conditions tested when compared to the wild type strain. In TSB media, after 72 h of culture, the presence of Caa or CaaB in Re2428 resulted in OD<sub>600nm</sub> of 2.96 ± 0.039 and 1.18 ± 0.009 respectively, both of which were much lower than growth of the wild type (OD<sub>600nm</sub> = 4.95 ± 0.35). In complementation strains, the localization of Caa in the cell during growth on TSB media did not present a significant difference (Figure 5B). In minimal media, however, a partial recovery of growth was observed only for Re2428/Caa (Figure 5C). These results indicate that overexpression of *caa* in Re2428 was incapable of recovering growth to



**Figure 4** Growth of the quadruple deletion mutant on TSB + 10% CO<sub>2</sub>. Cultures of *R. eutropha* H16 (□), *R. eutropha* CA quadruple deletion mutant Re2436 (H16  $\Delta$ can  $\Delta$ can2  $\Delta$ caa  $\Delta$ cag - ●) and the complemented strains, Re2436/pCan (○), Re2436/pCan2 (▼), Re2436/Caa (△) and Re2436/pCag (■) in TSB media in the presence of 10% CO<sub>2</sub>. Values represent average from three replicates with standard deviation as error bars.





wild type rates in either rich or minimal medium, even though CA activity was detected in cell extracts of Re2428/pCaa and Re2424/pCaaB (data not shown). The periplasmic localization of Caa, under the conditions tested, was only observed to affect growth of the cells to a significant extent when cultivated in minimal media.

#### Localization of Caa

In order to provide evidence for the periplasmic localization of Caa in *R. eutropha*, RFP was fused to the C-terminus of Caa to allow visualization of the localization in the cell.

RFP-tagged Caa and RFP alone (control) were expressed separately in the *R. eutropha* strain Re2061, which is unable to produce intracellular polyhydroxyalkanoate that may disturb the imaging. As shown in the fluorescent microscopy images presented in Additional file 1: Figure S4A, the fluorescence was concentrated near the outer perimeter of the cells with RFP-tagged Caa, as opposed to the RFP-only control, where fluorescence was diffused throughout the cytosol (Additional file 1: Figure S4B). This suggests that Caa enzymes are located near the outer perimeter of the cell, likely the periplasm.

#### Discussion

In this study, we evaluated four putative CA genes found in *R. eutropha* genome. The *can* and *can2* genes encode β-CA enzymes, the *caa* gene encodes a putative periplasmic α-CA, and the *cag* gene encodes a γ-like-CA/acetyltransferase. We demonstrated the ability of all four CA enzymes to catalyze the interconversion of CO<sub>2</sub> and bicarbonate, as purified enzymes and also when overexpressed in *R. eutropha*. We elucidated some of the complexities of CO<sub>2</sub> metabolism in *R. eutropha* and the evolutionary importance for the bacteria to retain four separate CA genes, which apparently present similar

activities in the cell, but which have unique roles within the machinery for controlling its CO<sub>2</sub> metabolism. In short, the presence of one enzyme does not exclude the importance of the others.

Previous microarray data produced in our laboratory demonstrated different expression behavior of the four CA genes when *R. eutropha* was growing on fructose or trioleate in the presence of high or low concentrations of nitrogen [31]. The *can* gene exhibited an expression decrease of 1.7 fold ( $p = 0.0048$ ) when cells entered into nitrogen depletion, *i.e.* the PHB production stage. On the other hand, *cag* exhibited an increase in expression of 3.5 fold ( $p = 0.011$ ) during the same growth condition. When changing the carbon sources from trioleate to fructose, *caa* exhibited an expression decrease of 3.0 fold ( $p = 0.001$ ). All of these previous observations support the idea that the four enzymes are not responsible for the same reactions in the cell, although Kusian et al. [1] proposed that, out of the four CA enzymes described in *R. eutropha*, only *can* plays an important role on CO<sub>2</sub> metabolism. Supporting our findings of independent roles of CAs in *R. eutropha*, [32-34] also reported and discussed the separate roles of different CAs expressed in the same organism. Dobrinski et al. [17] described four different CAs expressed at different levels depending on the physiological state of the microorganism in *T. crunogena*. Among other results, the observed differences in expression levels indicate a specific role of each CA enzyme in carbon fixation, pH homeostasis or other physiological activities not yet elucidated.

Overexpression of the CA enzymes separately in the wild type strain resulted in no observable difference in growth (Figure 2A) or PHB production (data not shown), but higher CA activities were detected in the cell extracts when compared to the wild type harboring the empty vector (Figure 2B). Examination of the purified CA enzymes (Figure 1), and CA overexpression

strains (Figure 2), showed that Caa exhibited greater activity using CO<sub>2</sub> as a carbon source. Under autotrophic growth conditions, when CO<sub>2</sub> is used as the sole carbon source, the growth and PHB accumulation of the CA overexpression strains presented significant differences when compared to each other and to the wild type strain (Figure 2C and D). One reason for the low growth and PHB production of the overexpression strains could be due to the altered regulation of gene expression.


The phenotype recovery effect of the overexpression of Caa in Re2428 (H16Δ*caa*) is more distinct when cells are grown in minimal media (Figure 5C), because in this environment, bicarbonate is required to synthesize essential amino acids and nucleotides, which are not supplied in the media [33].

We have identified and characterized an α-periplasmic CA, Caa, and demonstrated the importance of Caa for the transport of CO<sub>2</sub> and the supply of bicarbonate to *R. eutropha* cells during growth. Caa deletion, imbalanced expression, and mislocalization all disabled and adversely affected cell growth, presumably by severely impairing *R. eutropha* metabolism.

Several essential metabolic pathways require either CO<sub>2</sub> or bicarbonate as a substrate. Bicarbonate is the substrate of several important enzymes of central metabolism, such as phosphoenolpyruvate carboxylase, pyruvate carboxylase, acetyl-CoA carboxylase and methylcrotonyl CA, among others [3]. Calculated by Merlin et al. [33], the spontaneous diffusion of CO<sub>2</sub> in air and conversion to bicarbonate inside the cell are not sufficient for the metabolic needs of a bacterial cell. In

*R. eutropha* metabolism, bicarbonate is required for the elimination of a long lag phase during autotrophic growth [35], which is considered to be a negative feature for the use of this microorganism at industrial scale. CAII, a human α-CA, is known to play an important role in the transportation of CO<sub>2</sub> and supplying bicarbonate to mammalian cells [36] and in *T. crunogena*. Dobrinski et al. [17] also propose that an α-CA would be converting CO<sub>2</sub> to bicarbonate in the cells. In *R. eutropha*, Caa could be responsible for this same mechanism, since the equilibrium of Caa-catalyzed reaction lies towards the hydration of CO<sub>2</sub> to bicarbonate (Figure 1). Can, alternatively, could be responsible for the supplementation of CO<sub>2</sub>. These hypotheses were supported by the deletion of *caa* or *can* from the genome of *R. eutropha*. The lack of *can* resulted in a phenotype in which cells were dependent on external CO<sub>2</sub> supplementation in order to grow [1], this study. Without *caa* (i.e., strain Re2428), *R. eutropha* presented a similar phenotype as the *can* deletion strain, but the activities and substrate specificities of these two CA enzymes are remarkably different. Moreover, the growth phenotype of a *caa* deletion could not be compensated by the addition of CO<sub>2</sub> to the culture. These differences imply disparate physiological roles of both of these CA enzymes in the growth and maintenance of *R. eutropha*.

Additionally, both Caa and Can could not replace the roles of one another in *R. eutropha*, because a single deletion of either *can* or *caa* resulted in no growth under ambient CO<sub>2</sub> conditions [1], this study. Although *in vitro* activity suggested that the enzymes could have



complementary activities (Figure 1). Overall, we conclude in this study that Caa is unable to act as a replacement for Can, and vice versa. This conclusion is supported by the physical separation of both CAs in the cell, in which Caa is located in the periplasm and Can is believed to be in the cytosol.

The attempt to recover fully the phenotype of Re2428 by overexpressing a plasmid-borne *caa* gene was unsuccessful, as the overall growth yields of the complementary strains Re2428/pCaa and Re2428/pCaaB were much lower than that of the wild type, even when both versions of the enzyme, after purification, exhibited high activity (Figure 5A). This could be the result of an expression level imbalance of the plasmid-borne *caa* compared to that of wild type. Interestingly, the strain Re2428/pCaaB was unable to grow at the same rate as Re2428/pCaa in minimal media (Figure 5C), although both complemented strains grew similarly in TSB medium (Figure 5B). The growth difference could be a result of the differential localization of the two versions of Caa in the cell. The lack of a signaling peptide may have prevented the periplasmic positioning of Caa in the cell, and thus resulted in a lower bacterial growth. This effect can be more clearly observed in minimal media (Figure 5C). The results suggest that Caa is a periplasmic CA and its correct location in the cell is essential for the growth of *R. eutropha*. Overall, reintroduction of *caa* or *caaB* in Re2428 was able to recover only partial growth.

The isolated effect of the CAs in the genome could be observed after complementing the quadruple mutant, Re2436 (H16 $\Delta$ *can* $\Delta$ *caa* $\Delta$ *can2* $\Delta$ *cag*), with each separate enzyme expressed from a plasmid (Figure 4). In a background with no CA enzymes expressed, none of the individual CAs expressed *in trans* could fully recover the growth phenotype under all conditions tested. However, strain Re2436/pCan, expressing the Can enzyme, when grown in minimal media with fructose as the main carbon source could partially recover the growth phenotype (Figure 4). This reinforces the importance of having all four CAs in a complex system in order for carbon metabolism and cell growth to function properly.

The localization of Caa in the periplasm of the cell is crucial for the conversion of CO<sub>2</sub>, which is passively diffused through the membrane, to dissolved inorganic carbon (bicarbonate). The conversion of CO<sub>2</sub> to bicarbonate is important for the transportation of CO<sub>2</sub> into the cell, as CO<sub>2</sub> is extremely insoluble in aqueous solution and frequently diffuses in and out of the cell. Bicarbonate, on the other hand is negatively charged and highly soluble in aqueous solution, but poorly soluble in lipids. Transport of bicarbonate across the cell membrane must be assisted by CA [3,5]. Figure 6 depicts a diagram representing the role of CAs in *R. eutropha* cells. Can is likely the enzyme responsible for supporting

CO<sub>2</sub> fixation reaction by providing CO<sub>2</sub> to RuBisCO in the cytosol. The same reaction can potentially be performed by Cag, but its role is elusive and thus merits further investigation. Caa is responsible for sequestering the CO<sub>2</sub> diffused into the cell and converting it to bicarbonate for cellular metabolism, while Can2 has a possible role in pH homeostasis. The current work provides a springboard to understanding the differential roles of each CA enzyme in the physiology and CO<sub>2</sub> homeostasis of *R. eutropha*. The exact interplay of each of these enzymes in the cell still continues to be an active area of study.

## Additional file

**Additional file 1: Figure S1.** Growth of complemented deletion strains. **Figure S2.** Light microscopy of deletion strains. **Figure S3.** Growth of H16 and Re2427 with different initial pH values. **Figure S4.** Fluorescent microscopy of Caa\_RFP fusion protein expressed in Re2061. **Table S1.** Oligonucleotide primers used in this study [37-43].

## Competing interest

The authors declare that there is no conflict of interest.

## Acknowledgements

The authors thank Mr. John W. Quimby for the critical review of this manuscript, and Professor Pamela Silver and Dr. Buz Barstow from the Department of Systems Biology at Harvard Medical School for their help with the autotrophic growth experiments. This work is funded by the US Department of Energy, Advanced Research Projects Agency—Energy (ARPA-E). We thank our ARPA-E collaborators Dr. Mark Worden and Ms. Yangmu Liu for their helpful discussions and support throughout the course of this study.

## Author details

<sup>1</sup>Department of Biology, Massachusetts Institute of Technology, Bldg. 68-370, Cambridge, MA 02139, USA. <sup>2</sup>Department of Chemistry, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA. <sup>3</sup>Department of Bioengineering, University of Massachusetts Dartmouth, 285 Old Westport Road, North Dartmouth, MA 02747, USA. <sup>4</sup>Division of Health Sciences and Technology, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Bldg. 68-370, Cambridge, MA 02139, USA. <sup>5</sup>Engineering Systems Division, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Bldg. 68-370, Cambridge, MA 02139, USA.

Received: 17 December 2013 Accepted: 23 December 2013

Published: 10 January 2014

## References

1. Kusian B, Sültemeyer D, Bowien B (2002) Carbonic anhydrase is essential for growth of *Ralstonia eutropha* at ambient CO<sub>2</sub> concentrations. *J Bacteriol* 184:5018–5026
2. Lionetto MG, Caricato R, Giordano ME, Erroi E, Schettino T (2012) Carbonic Anhydrase and Heavy Metals, Biochemistry. In: Deniz E (ed) Tech., <http://www.intechopen.com/books/biochemistry/carbonic-anhydrase-and-heavy-metals>. ISBN: 978-953-51-0076-8
3. Smith KS, Ferry JG (2000) Prokaryotic carbonic anhydrases. *FEMS Microbiol Rev* 24:335–366
4. Fasseas MK, Tsikou D, Fletmetakis E, Katinakis P (2011) Molecular and biochemical analysis of the a class carbonic anhydrases in *Caenorhabditis elegans*. *Mol Biol Rep* 38:1777–1785
5. Marcus EA, Moshfegh AP, Sachs G, Scott DR (2005) The periplasmic  $\alpha$ -carbonic anhydrase activity of *Helicobacter pylori* is essential for acid acclimation. *J Bacteriol* 187:729–738
6. Tripp BC, Smith K, Ferry JG (2001) Carbonic anhydrase: new insights for an ancient enzyme. *J Biol Chem* 28:48615–48618

7. Supuran CT (2004) Carbonic Anhydrases: Catalytic and Inhibition Mechanisms, Distribution and Physiological Roles. In: Supuran CT, Scozzafava A, Conway J (eds) Carbonic anhydrase: its inhibitors and activators. CRC press LLC, Boca Raton, USA, p 376
8. Brigham CJ, Zhila N, Shishatskay E, Volova TG, Sinskey AJ (2012b) Manipulation of *Ralstonia eutropha* carbon storage pathways to produce useful bio-based products. In: Wang (ed) Reprogramming microbial metabolic pathways X. Springer Science+Business Media Dordrecht. Subcellular Biochemistry 64, DOI 10.1007/978-94-007-5055-5\_17
9. Budde CF, Riedel SL, Willis LB, Rha C, Sinskey AJ (2011) Production of poly(3-hydroxybutyrate-co-3-hydroxyhexanoate) from plant oil by engineered *Ralstonia eutropha* strains. *Appl Environ Microbiol* 77:2847–2854
10. Li H, Oppenorth PH, Wernick DG, Rogers S, Wu TY, Higashide W, Malati P, Huo YX, Cho KM, Liao JC (2012) Integrated electromicrobial conversion of CO<sub>2</sub> to higher alcohols. *Science* 335:1596
11. Lu J, Brigham CJ, Gai CS, Sinskey AJ (2012) Studies on the production of branched-chain alcohols in engineered *Ralstonia eutropha*. *Appl Microbiol Biotechnol* 96:283–297
12. Kusian B, Bowien B (1997) Organization and regulation of *cbb* CO<sub>2</sub> assimilation genes in autotrophic bacteria. *FEMS Microbiol Rev* 21:135–155
13. Codd GA, Kuenen JG (1987) Physiology and biochemistry of autotrophic bacteria. *Antonie Van Leeuwenhoek* 53:3–14
14. Badger MR, Price GD (2003) CO<sub>2</sub> concentrating mechanisms in cyanobacteria: molecular components, their diversity and evolution. *J Exp Botany* 54:609–622
15. Bonacci W, Teng PK, Afonso B, Niederholtmeyer H, Grob P, Silver PA, Savage DF (2012) Modularity of a carbon-fixing protein organelle. *Proc Natl Acad Sci USA* 109:478–483
16. Pohlmann A, Fricke WF, Reinecke F, Kusian B, Liesegang H, Cramm R, Eitinger T, Ewering C, Pötter M, Schwartz E, Strittmatter A, Voß I, Gottschalk G, Steinbüchel A, Friedrich B, Bowien B (2006) Genome sequence of the bioplastic-producing “Knallgas” bacterium *Ralstonia eutropha* H16. *Nat Biotechnol* 24:1257–1262
17. Dobrinski KP, Boller AJ, Scott KM (2010) Expression and function of four carbonic anhydrase homologous in the deep-sea chemolithoautotroph *Thiomicrospira crunogena*. *Appl Environ Microbiol* 76:3561–3567
18. Simon R, Priefer U, Pühler A (1983) A broad host range mobilization system for *in vivo* genetic engineering: transposon mutagenesis in gram-negative bacteria. *Nat Biotechnol* 1:784–791
19. Kovach ME, Elzer PH, Hill DS, Robertson GT, Farris MA, Roop RM 2nd, Peterson KM (1995) Four new derivatives of the broad-host-range cloning vector pBRR1MCS, carrying different antibiotic-resistance cassettes. *Gene* 166:175–176
20. Novy R, Drott D, Yaeger K, Mierendorf R (2001) Overcoming the codon bias of *E. coli* for enhanced protein expression. *iNovations* 12:1–3
21. Bertani G (1951) The mode of phage liberation by lysogenic *Escherichia coli*. *J Bacteriol* 62:293–300
22. Leadbeater L, Bowien B (1984) Control of autotrophic carbon assimilation in *alcaligenes eutrophus* by inactivation and reactivation of phosphoribulokinase. *J Bacteriol* 157:95–99
23. Sambrook J, Russell DW (2001) Molecular cloning: a laboratory manual, 3rd edn. Cold Spring Harbor Laboratory Press, Cold Spring, USA
24. Slater S, Houmiel KL, Tran M, Mitsky TA, Taylor NB, Padgett SR, Gruys KJ (1998) Multiple  $\beta$ -ketothiolases mediate poly(beta-hydroxyalkanoate) copolymer synthesis in *Ralstonia eutropha*. *J Bacteriol* 180:1979–1987
25. Petersen TN, Brunak S, von Heijne G, Nielsen H (2011) SignalP 4.0: discriminating signal peptides from transmembrane regions. *Nat Methods* 8:785–786
26. Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA 3rd, Smith HO (2009) Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat Methods* 6:343–345
27. Sundaram V, Rumbolo P, Grubb J, Strisciuglio P, Sly WS (1986) Carbonic anhydrase II deficiency: diagnosis and carrier detection using differential enzyme inhibition and inactivation. *Am J Hum Genet* 38:125–136
28. Bradford MM (1976) Rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248–254
29. Karr DB, Waters JK, Emerich DW (1983) Analysis of poly- $\beta$ -hydroxybutyrate in *Rhizobium japonicum* Bacteroids by ionexclusion high-pressure liquid chromatography and UV detection. *Appl Environ Microbiol* 46:1339–1344
30. York GM, Lupberger J, Tian J, Lawrence AG, Stubbe J, Sinskey AJ (2003) *Ralstonia eutropha* H16 encodes two and possibly three intracellular poly(D-3-hydroxybutyrate) depolymerase genes. *J Bacteriol* 185:3788–3794
31. Brigham CJ, Speth DR, Rha C, Sinskey AJ (2012) Whole-genome microarray and gene deletion studies reveal regulation of the polyhydroxyalkanoate production cycle by the stringent response in *Ralstonia eutropha* H16. *Appl Environ Microb* 78:8033–8044
32. Kupriyanova E, Villarejo A, Markelova A, Gerasimenko L, Zavarzin G, Samuelsson G, Los DA, Pronina N (2007) Extracellular carbonic anhydrases of the stromatolite-forming cyanobacterium *Microcoleus chthonoplastes*. *Microbiol* 153:1149–1156
33. Merlin C, Masters M, McAteer S, Coulson A (2003) Why is carbonic anhydrase essential to *Escherichia coli*? *J Bacteriol* 185:6415–6424
34. Smith KS, Jakubczik C, Whittam TS, Ferry JG (1999) Carbonic anhydrase is an ancient enzyme widespread in prokaryotes. *Proc Natl Acad Sci USA* 96:15184–15189
35. Repaske R, Ambrose CA, Repaske AC, De Lacy ML (1971) Bicarbonate requirement for elimination of the lag period of *Hydrogenomonas eutropha*. *J Bacteriol* 107:712–717
36. Sterling D, Reithmeier RA, Casey JR (2001) Carbonic anhydrase: in the driver's seat for bicarbonate transport. *JOP* 4:165–170
37. Amoroso G, Morell-Avrahov L, Müller D, Klug K, Sültemeyer D (2005) The gene NCE103 (YNL036w) from *Saccharomyces cerevisiae* encodes a functional carbonic anhydrase and its transcription is regulated by the concentration of inorganic carbon in the medium. *Molec Microbiol* 56:549–558
38. Chirica LC, Elleby B, Jonsson BH, Lindskog S (1997) The complete sequence, expression in *Escherichia coli*, purification and some properties of carbonic anhydrase from *Neisseria gonorrhoeae*. *Eur J Biochem* 244:755–760
39. Kupriyanova EV, Sinetova MA, Markelova AG, Allakhverdiev SI, Los DA, Pronina NA (2011) Extracellular  $\beta$ -class carbonic anhydrase of the alkaliphilic cyanobacterium *Microcoleus chthonoplastes*. *J Photochem Photobiol B* 103:78–86
40. Luca VD, Vullo D, Scozzafava A, Carginale V, Rossi M, Supuran CT, Capasso C (2013) An  $\alpha$ -carbonic anhydrase from the thermophilic bacterium *Sulphurihydrogenibium azorense* is the fastest enzyme known for the CO<sub>2</sub> hydration reaction. *Bioorg Med Chem* 21:1465–1469
41. Mogensen EG, Janbon G, Chaloupka J, Steegborn C, Fu MS, Moyrand F, Klengel T, Pearson DS, Geeves MA, Buck J, Levin LR, Mühlischlegel FA (2006) *Cryptococcus neoformans* senses CO<sub>2</sub> through the carbonic anhydrase Can2 and the adenyl cyclase Cac1. *Eukariotic Cell* 5:103–111
42. Sültemeyer D (1998) Carbonic anhydrase in eukaryotic algae: characterization, regulation, and possible function during photosynthesis. *Can J Bot* 76:962–972
43. Wilbur KM, Anderson NG (1948) Electrometric and colorimetric determination of carbonic anhydrase. *J Biol Chem* 176:147–154

doi:10.1186/2191-0855-4-2

**Cite this article as:** Gai et al.: Insights into bacterial CO<sub>2</sub> metabolism revealed by the characterization of four carbonic anhydrases in *Ralstonia eutropha* H16. *AMB Express* 2014 4:2.

**Submit your manuscript to a SpringerOpen® journal and benefit from:**

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)