

ORIGINAL ARTICLE

Open Access



Microbicidal activity of N-chlorotaurine in combination with hydrogen peroxide

Jasmin Mustedanagic¹, Valdecir Farias Ximenes² and Markus Nagl^{1,3*} 

Abstract

N-chlorotaurine (NCT) and hydrogen peroxide are powerful endogenous antiseptics. In vivo, the reaction between hydrogen peroxide and metal ions leads to the formation of free hydroxyl radicals, which have an increased bactericidal activity. This study examined whether there is an additive antimicrobial effect of NCT combined with hydrogen peroxide. Additionally, it was tested if the additive effect is based on the formation of free radicals. We found by luminometry that, in the presence of H₂O₂, NCT caused a slow and long-lasting production of singlet oxygen in contrast to HOCl, where this burst occurred instantaneously. Both NCT and hydrogen peroxide (1.0 and 0.1%) demonstrated bactericidal and fungicidal activity. At pH 7.1 and 37 °C, hydrogen peroxide (1%, 294 mM) showed a stronger bactericidal and particularly fungicidal activity than NCT (1%, 55 mM), whereas at pH 4.0 and also in the presence of 5.0% peptone NCT revealed a stronger bactericidal activity. A combination of NCT and hydrogen peroxide led to an increased bactericidal but no increased fungicidal activity compared to both substances alone. The additive effect against bacteria was not removed in the presence of the radical scavengers NaN₃, DMSO, or peptone. As a conclusion, NCT and hydrogen peroxide used concurrently interact additive against a range of microorganisms. However, the results of this study suggest that the additive effect of NCT combined with hydrogen peroxide is rather not based on the formation of free radicals.

Keywords: *N*-Chlorotaurine, Hydrogen peroxide, Singlet oxygen, Microbicidal, Antimicrobial agent, Oxidant

Introduction

Human phagocytes are activated by invading microorganisms and produce several reactive oxygen species (ROS) by an enzymatic cascade called oxidative burst (Klebanoff 1968). Superoxide (O₂⁻) and hydrogen peroxide (H₂O₂) are products of NADPH oxidase and superoxide dismutase, respectively, and hypochlorous acid (HOCl) is formed from H₂O₂ and chloride by myeloperoxidase (Weiss et al. 1982; Zgliczynski et al. 1971). HOCl immediately reacts among others with amino groups to create chloramines (R-NHCl), also designated as long-lived oxidants because of their lower reactivity. *N*-chlorotaurine (Cl-HN-CH₂-CH₂-SO₃⁻, NCT) is the most abundant representative of this class of compounds (Grisham et al. 1984). All these oxidants are thought to be


involved in the killing of microorganisms during inflammation (Klebanoff et al. 2013). However, their spectrum of functions is broader, such as signal transduction by H₂O₂ (Kim et al. 2002; Mongkolsuk and Helmann 2002), or anti-inflammatory effects by NCT (Kim and Cha 2014; Marcinkiewicz and Kontny 2014). Moreover, chemically synthesized H₂O₂ and HOCl are in use as antiseptics in human medicine (Bruch 2007; Wilkins and Unverdorben 2013), and NCT is particularly suited for treatment of infections of sensitive body sites according to several studies (Gottardi and Nagl 2010; Nagl et al. 2003; Neher et al. 2004; Teuchner et al. 2005).

As an interesting aspect, the mentioned oxidants may also react among one another, whereby further microbicidal ROS are formed. The reaction between HOCl and H₂O₂ to form singlet oxygen (¹O₂) (Eq. 1) was discovered by Khan and Kasha in (1963), and it was studied in details due to its physiological and pathological relevance, including its involvement in signaling and microbicidal

*Correspondence: m.nagl@i-med.ac.at

³ Division of Hygiene and Medical Microbiology, Medical University of Innsbruck, Schöpfstr. 41, 1st Floor, 6020 Innsbruck, Austria
Full list of author information is available at the end of the article

functions (Khan and Kasha 1963; Steinbeck et al. 1992; Stief and Fareed 2000).



The oxidation of H_2O_2 leading to ${}^1\text{O}_2$ is not a property exclusive of HOCl (Miyamoto et al. 2014), but also shared by other halogenating species as hypobromous acid (HOBr) (Kanofsky et al. 1988), chloramines, including NCT (Stief 2003) and N-bromotaurine (De Carvalho et al. 2016). However, although the reactivity of NCT with H_2O_2 leading to ${}^1\text{O}_2$ has been proposed (Stief et al. 2001; Stief 2003), as far as we know, it was never investigated in details. Here we identified an advantage in the use of NCT compared to HOCl regarding the production of ${}^1\text{O}_2$.

The aim of this study was to characterize the reaction of NCT and hydrogen peroxide with production of singlet oxygen and to investigate a possible additive bactericidal or fungicidal effect of both compounds.

Materials and methods

Chemicals

Taurine, melatonin, hydrogen peroxide and deuterium oxide were purchased from Sigma-Aldrich Chemical Co. (St. Louis, MO, USA). Working solution of hypochlorous acid (HOCl) 100 mM was prepared by diluting a 5% commercial solution in water. The concentration of HOCl was determined spectrophotometrically after diluting the working solution in 0.01 M NaOH, pH 12 ($\lambda_{\text{max}} = 292 \text{ nm}$, $\epsilon = 350/\text{M cm}$). N-Chlorotaurine (NCT) (molecular weight 181.57 g/mol, lot 2015-02-05) was prepared as crystalline sodium salt in our laboratory (M. Nagl, Innsbruck) at pharmaceutical grade, as reported (Gottardi and Nagl 2002), stored at minus 20 °C, and freshly dissolved in sterile 0.1 mM sodium phosphate buffer at pH 7.1 to a concentration of 55 mM (1%), 5.5 mM (0.1%), 1 mM (0.018%) or 0.25 mM (0.0045%) for each experiment.

Hydrogen peroxide (H_2O_2) 100 mM was prepared by dilution of a 30% commercial solution (Merck, Darmstadt, Germany) in water. The concentration of H_2O_2 was determined spectrophotometrically by its absorbance ($\lambda_{\text{max}} = 240 \text{ nm}$, $\epsilon = 43.6/\text{M cm}$). The chemicals used for preparation of phosphate buffer and solutions were of analytical grade. Ultrapure Milli-Q water (Millipore, Belford, MA, USA) was used for the preparation of buffers and solutions. Bacto™ peptone from Becton–Dickinson and Company (NJ, USA) was dissolved in distilled water to a 10% stock solution and autoclaved. Sodium azide (NaN_3) was dissolved in distilled water to a 0.65% (100 mM) stock, dimethyl sulfoxide (DMSO) to a 10% stock (1.28 M) in 0.1 M phosphate buffer. Catalase from *Micrococcus lysodeikticus* containing 65,000–150,000 U/

ml was from Sigma-Aldrich (Germany). Chloramine T from Merck was dissolved in 0.1 M phosphate buffer to 0.005% (0.178 mM).

Studies of the reaction between NCT or HOCl and H_2O_2

Consumption of NCT and HOCl were monitored by their absorbances at 252 and 290 nm, respectively, using a Perkin Elmer Lambda 25 UV–visible spectrophotometer (Shelton, CT, USA). The reaction mixtures were composed of 0.25 mM NCT or HOCl, and 0.5 mM H_2O_2 in 0.1 M phosphate buffer, pH 7.1 and 37 °C. The production of ${}^1\text{O}_2$ was monitored by its dimol light emission or, indirectly, by the light emission generated by its reaction with melatonin using a plate luminometer (Centro Microplate Luminometer LB960, Berthold Technologies, Oak Ridge, TN, USA). The reaction mixtures were composed of 1.0 mM NCT or HOCl, 20 mM H_2O_2 , in the presence or absence of 1 mM melatonin in 0.1 M phosphate buffer, pH 7.1 and 37 °C. The reactions were triggered by the addition of H_2O_2 .

Bacteria and fungi

Bacteria and yeasts deep frozen for storage were grown on Mueller–Hinton agar plates (Oxoid, Hampshire, UK) and subcultivated overnight in tryptic soy broth (Merck) at 37 °C. Subsequently, they were washed twice in 0.9% saline before use. Strains used were *Staphylococcus aureus* ATCC 25923 and 6538, *Pseudomonas aeruginosa* ATCC 27853, *Escherichia coli* ATCC 11229, and *Candida albicans* CBS 5982 (60% pseudohyphae and 40% blastoconidia). *Aspergillus fumigatus* ATCC 204305 was grown on Sabouraud agar (Becton & Dickinson, Heidelberg, Germany) for 72 h. Suspensions of conidia were gained by harvesting them from the agar plates with 5.0 ml of 0.9% saline plus 0.01% Tween 20, followed by 10- μm filtration (CellTrics; Partec GmbH, Görlitz, Germany) to gain a pure conidia suspension without hyphae and three washing steps in phosphate-buffered saline (Lackner et al. 2015).

Time-kill assays (Lackner et al. 2015; Martini et al. 2012)

All experiments were done at 37 °C in a water bath. NCT (1.98 ml) was mixed with H_2O_2 (1.98 ml) in 0.1 M phosphate buffer (pH 7.1) or 0.1 M sodium acetate buffer (pH 4.0) to final concentrations of 1% (equals 55 mM NCT and 294 mM H_2O_2) or 0.1% each. In parallel, 3.96 ml of 1 or 0.1% NCT and 3.96 ml of 1 or 0.1% H_2O_2 were investigated in each experiment. A control in phosphate or acetate buffer without additives was done in parallel. In separate experiments, single radical scavengers were added to all test tubes. Thereby, respective volumes of the stock solutions were mixed immediately before the start of the test, for instance 1.98 ml of 2% NCT plus 2% H_2O_2

with 1.98 ml 10% peptone. Final concentrations were 5% peptone, 5% DMSO, or 10 mM sodium azide. To 3.96 ml of the test solutions, 40 μ l of the respective suspension containing bacteria or fungi were added at time zero and vortexed. Final starting concentrations of microorganisms were $1.6\text{--}6.6 \times 10^6$ colony forming units (cfu)/ml for *S. aureus* ATCC 25923, $1.1\text{--}2.3 \times 10^7$ cfu/ml for *S. aureus* ATCC 6538, $1.1\text{--}3.1 \times 10^7$ cfu/ml for *P. aeruginosa* and *E. coli*, $2.0\text{--}4.4 \times 10^5$ cfu/ml for *C. albicans*, $4.4\text{--}1.0 \times 10^6$ cfu/ml for *A. fumigatus*. Incubation times ranged between 1 and 240 min and are indicated in the figures. At the end of each incubation time, aliquots of 100 μ l were diluted in 900 μ l NCT-inactivating solution consisting of 890 μ l of 3% sodium thiosulphate plus 10 μ l (approximately 1000 U) catalase in distilled water. Aliquots of 50 μ l of this solution were spread on Mueller–Hinton agar plates in duplicate using an automatic spiral plater (model WASP 2, Don Whitley, Shipley, United Kingdom). The detection limit was 100 cfu/ml, taking into account both plates and the previous 10-fold dilution in the inactivating solution. Plates were grown for 48 h (bacteria) to 72 h (fungi) at 37 °C, and the number of cfu was counted. Plates with no growth or only a low cfu count were grown for up to 5 days (bacteria, *Candida*) or ten days (*Aspergillus*). Controls, i.e. plain 0.1 M phosphate buffer with and without scavengers (peptone, DMSO, NaN_3) were performed in parallel. Inactivation controls, where NCT and H_2O_2 were mixed with their inactivators thiosulphate plus catalase immediately before addition of pathogens at low cfu counts, showed full survival of bacteria and fungi. This proved rapid and sufficient inactivation.

Time-kill assays with sequential treatment of NCT and H_2O_2

To investigate if there is an additive bactericidal effect after sequential incubation in both compounds, the pellet of washed bacteria (*S. aureus* ATCC 6538, *E. coli*, *P. aeruginosa*) was resuspended in the slower acting agent, 1% NCT, in 0.1 M phosphate buffer (pH 7.1) to $2\text{--}5 \times 10^9$ cfu/ml at room temperature first. The incubation time was 1 min. Controls were treated with phosphate buffer without NCT. After that time, the cfu count is not influenced, but the surface of the bacteria becomes chlorinated (“chlorine cover”) (Gottardi and Nagl 2005). Then, the bacteria were centrifuged at $4000 \times g$ for 5 min, washed in 0.9% NaCl, centrifuged again, and resuspended in saline. Subsequently, 40 μ l of the bacterial suspension was added to 3.96 ml 1% H_2O_2 (0.3% H_2O_2 for tests with *E. coli* and *P. aeruginosa*) in 0.1 M phosphate buffer (pH 7.1), and incubated at 37 °C (controls in buffer without H_2O_2). Quantitative cultures from aliquots after different incubation times were performed as described in the previous paragraph.

Statistics


The data are presented as mean values and standard deviations (SD) of at least three independent experiments. Student’s unpaired *t* test in case of two groups or one-way analysis of variance (ANOVA) and Tukey’s multiple-comparison test in case of more than two groups were used to test for a difference between the test and control groups. A *P* value of <0.05 was considered significant for all tests. Calculations were done with the GraphPad Prism 6.01 software (GraphPad, Inc., La Jolla, CA, USA).

To gain an improved survey on the microbicidal activity of NCT against the different strains, the recently introduced Integral Method was used, which transforms the whole killing curve (\log_{10} cfu/ml versus time) into one value of “bactericidal activity (BA)” (Gottardi et al. 2015). The higher the value, the stronger is the microbicidal activity. Moreover, the method allows an expanded statistical analysis with the tests mentioned above, particularly between killing curves with small differences.

Results

Reaction between NCT or HOCl and H_2O_2



We found that, differently of HOCl, which produces an instantaneous burst of $^1\text{O}_2$ in the presence of H_2O_2 , NCT caused a slow and long-lasting production of this electronically excited form of molecular oxygen. Figure 1 shows the consumption of HOCl and NCT provoked by the addition of H_2O_2 . While HOCl was almost completely depleted in less than 5 s, NCT lost less than 2% of its initial concentration in 20 min. The reactions were also monitored by the weak chemiluminescence generated by dimol emission of $^1\text{O}_2$ (Lengfelder et al. 1983). Figure 2a shows that, although of low intensity and of a short period (less than 10 s), chemiluminescence was detected by the reaction between HOCl and H_2O_2 . On the other hand, due to its lower reactivity, light emission was not detected using NCT. Hence, to improve the efficiency, we added melatonin, which promptly reacts with $^1\text{O}_2$ leading to light emission through the formation of an unstable dioxetane intermediate (Lu et al. 2002; Matuszak et al. 2003). The result in Fig. 2b shows that the emission of light for the reaction between HOCl and H_2O_2 was increased about 3 orders of magnitude in the presence of melatonin, reinforcing the involvement of $^1\text{O}_2$. Finally, Fig. 3 shows the kinetic profile of light emission when HOCl was substituted by NCT in the presence of melatonin and H_2O_2 . Corroborating with the results obtained by monitoring the consumption of NCT, the production of $^1\text{O}_2$ took place in more than 30 min, which is an evidence of the slow, but efficient reactivity of this chloramine with H_2O_2 . Of note, according controls disclosed that light emission was not due to direct reaction of melatonin with H_2O_2 or with NCT (not showed). We also found that the reaction



rate was still lower at alkaline pH, which is consistent with the lower reactivity of NCT in this condition. Additional evidence of the formation of ¹O₂ was obtained by addition of deuterated water in the reaction medium, which increase its lifetime (Kim et al. 2016) and, consequently, the efficiency of the reaction (Fig. 3).

Microbicidal activity of NCT and H₂O₂

Against all bacterial test strains, the combination of NCT and H₂O₂ showed a more rapid killing than the single compounds. In phosphate buffer at pH 7.1, H₂O₂ had a throughout stronger activity than NCT. This is illustrated in Fig. 4. To investigate the influence of singlet oxygen formed by NCT plus H₂O₂, sodium azide as a scavenger was added to the test solutions at a concentration of 10 mM, which is a threshold that just does not kill bacteria (Sabbahi et al. 2008). Surprisingly, the significantly more rapid killing by the combination was still present (Fig. 5). The same was true if 5% DMSO, a hydroxyl radical scavenger, was added (data not shown). Finally, we added 5% peptone as a general quencher of oxidants (Fig. 6). Of note, NCT plus H₂O₂ was still the highly significantly strongest bactericidal solution. In addition, the already known enhancing effect of organic matter on







Fig. 4 Bactericidal activity of 1% NCT (filled square), 1% H_2O_2 (filled triangle), and 1% NCT plus 1% H_2O_2 (inverted triangle) at pH 7.1 and 37 °C. Control in phosphate buffer without additives (filled circle). Mean values \pm SD of three to four independent experiments. **P < 0.01 versus all other values. BA "bactericidal activity" [\log_{10} cfu/min] as a quantitative measure for the strength of killing calculated by the integral method for the whole killing curve according to (Gottardi et al. 2015). The higher the value, the higher the microbicidal activity

NCT was seen again (Gottardi et al. 2014), while H_2O_2 was weakened. Therefore, NCT turned out to kill rather more rapidly than H_2O_2 under these conditions.

A similar result was seen at pH 4 in acetate buffer where 0.1% NCT and 0.1% H_2O_2 were tested. Again, the combination showed the strongest effect followed by NCT and H_2O_2 (BA = $0.98 \pm 0.15 \log_{10}$ cfu/min for

NCT plus H_2O_2 , 0.60 ± 0.07 for NCT, and 0.25 ± 0.02 for H_2O_2).

Against *C. albicans* and *A. fumigatus* in phosphate buffer, no difference was found between NCT plus H_2O_2 and H_2O_2 alone (Fig. 7), while NCT killed fungi much slower than bacteria as expected from former work (Nagl et al. 2001).



In some experiments, 0.005% CAT was used instead of 1% NCT against *S. aureus* 6538. The BA value of 1% H₂O₂ was 0.28 ± 0.02 , that of CAT 0.55 ± 0.05 and that of CAT plus 1% H₂O₂ 0.67 ± 0.07 log₁₀ cfu/min. The difference between CAT and CAT plus H₂O₂ did not reach significance but a trend after 3 independent experiments (*P* = 0.0676).


Time-kill assays with sequential treatment of NCT and H₂O₂

When NCT-pretreated, chlorine-covered bacteria were incubated in H₂O₂, a differential result was gained, depending on the species used. With *S. aureus*, the chlorine cover did not enhance the susceptibility to H₂O₂ (Fig. 8). By contrast, *E. coli* and *P. aeruginosa* were killed significantly more rapidly by H₂O₂ if pretreated with NCT (Fig. 8) (*P* < 0.01).

Discussion

Both hydrogen peroxide and NCT are important low reactive components of the oxidative armament of human granulocytes and monocytes and can be used as endogenous antiseptics in human medicine (Baldry 1983; Gottardi et al. 2013; Gottardi and Nagl 2010; Winterbourn and Kettle 2013). For both regards, investigations

of the interaction between these molecules may be of interest and contribute to elucidate biological processes. A slow and long-lasting production of singlet oxygen by NCT plus H₂O₂ was characterized for the first time in this study. Concentrations of the oxidants were adjusted to reveal good quantitative results. Two procedures were used to monitor the reactions and to highlight the differences between NCT and its precursor HOCl. In the first one, the higher reactivity of HOCl compared to NCT was clearly demonstrated by its rate of consumption. However, the concomitant production of singlet oxygen, monitored by its phosphorescence emission, is uneasy to follow due to the low efficiency of light emission. This was the reason to add melatonin in the medium, since its reactivity with singlet oxygen is well-known and characterized by the efficient emission of chemiluminescence (Lu et al. 2002). Scheme 1 depicts a mechanistic proposal for production of singlet oxygen and the emission of chemiluminescence through generation of a dioxetane intermediate. It is worth of note that there are other oxidative pathways by which melatonin could generate light emission, however, this was not the case using NCT or H₂O₂ alone. In addition, the amplification of the light emission provoked by deuterium oxide is an additional evidence of the intermediate singlet oxygen.




in 5% peptone: NCT + $H_2O_2 > NCT \geq H_2O_2$

Fig. 6 Bactericidal activity of 1% NCT (filled square), 1% H_2O_2 (filled triangle), and 1% NCT plus 1% H_2O_2 (inverted triangle) at pH 7.1 and 37 °C in the presence of 5% peptone. Control in phosphate buffer plus 5% peptone (filled circle). Mean values \pm SD of three to four independent experiments. $**P < 0.01$ versus all other values. BA values calculated as in Fig. 4

We hypothesized that production of 1O_2 would increase the microbicidal activity of NCT and H_2O_2 if they were used in combination. For these investigations, we applied

higher, clinically applied concentrations and gained reasonable killing times of microorganisms. Actually, against bacteria an additive effect of both compounds was found.



It was not strong, but reached high significance compared to the single components. This was true for single incubation time points as well as for the whole killing curves condensed by the integral method (Gottardi et al. 2015), which proved to be of high advantage for comparison of curves. The additive effect was independent of the pH and still present at pH 4. In acidic environment, both single NCT and H_2O_2 showed stronger killing than at pH 7, but this was much more pronounced with NCT. Therefore, NCT exerted a stronger bactericidal effect than H_2O_2 at pH 4. Active chlorine compounds typically and markedly increase their microbicidal activity at acidic pH due to formation of further oxidizing species and due to loss of negative charges of the bacterial surface [for details see (Gottardi et al. 2013)].

We further hypothesized that the additive effect of NCT and H_2O_2 would disappear in the presence of scavengers of oxygen radicals (DMSO, sodium azide). However, this assumption turned out as not true. The combination was still stronger bactericidal, which was further confirmed in the presence of 5% peptone, a general scavenger of oxidants. These results indicate that formed singlet oxygen might not be responsible for this effect. A possible explanation is a combined attack of NCT and H_2O_2 on the bacterial cell wall and

cell membrane. This may lead to earlier penetration into the microorganism followed by its rapid irreversible inactivation. In this regard, however, we cannot completely discard the involvement of singlet oxygen, since it is possible that azide (N_3^-), a charged molecule, and the other scavengers had not total access to the site of generation of singlet oxygen in the cell covers of the bacteria.

Sequential application of NCT and H_2O_2 , revealing an additive killing effect in Gram-negative bacteria but not in *S. aureus*, rather confirms that singlet oxygen is not responsible. Combined attack and more rapid destruction of the bacterial covers followed by slightly more rapid penetration of oxidation capacity into the bacteria appears to be a conclusive hypothesis in our opinion. Detailed contribution of the multitude of single reaction partners/products are unknown, although the basic chemical reactions have been elucidated (Gottardi et al. 2013; Peskin et al. 2009). From previous studies it is known that the chlorine cover attached by sublethal treatment with NCT does not kill the microorganisms but removes their virulence and causes a lag of regrowth and postantibiotic effect (Gottardi and Nagl 2005; Lackner et al. 2015; Nagl et al. 1999). Therefore, we think that the damage of the surface by the chlorine cover





Fig. 8 Bactericidal activity of H₂O₂ (filled triangle) and of H₂O₂ against chlorine-covered bacteria that were pretreated with 1% NCT for 1 min (inverted triangle). 1% H₂O₂ against *S. aureus* ATCC 6538, 0.3% H₂O₂ against *E. coli* and *P. aeruginosa*. Controls in phosphate buffer without additives, without (filled circle) or with chlorine cover (Asterisk). Mean values ± SD of three to four independent experiments. **P < 0.01 versus all other values. BA values calculated as in Fig. 4

promotes the attack by H₂O₂ at least in Gram-negatives. The thicker Gram-positive cell wall seems to resist few min longer to penetration of the used oxidants than the Gram-negative one.

The relatively slow killing of bacteria and fungi by millimolar NCT appears to approximately correlate with its penetration into the cytosol of these organisms. Slow penetration of 1 mM NCT into endothelial cells has been

found (Peskin et al. 2005), which was confirmed in our laboratory with 55 mM NCT using keratinocytes (A431), lung epithelial cells (A549), and *Aspergillus fumigatus* (M. Nagl, A. Windisch, unpublished results).

Against fungi, no additive effect was found, at least at the tested concentration. The activity of NCT against fungi is markedly lower than that of H₂O₂ in phosphate buffer, probably due to slow penetration (Nagl et al. 2001,



2002). Obviously, the combined attack is not sufficiently strong to kill fungi more rapidly.

Under protein load, H_2O_2 showed the expected decrease of activity, while NCT was markedly increased. The latter can be explained by transchlorination from NCT to amino groups of peptone, whereby among others low molecular weight chloramines are formed in equilibrium, which have stronger microbicidal activity (Gottardi et al. 2014). This is particularly true for monochloramine (NH_2Cl) because of its higher lipophilicity (Gottardi et al. 2007). The enhancement of activity under protein load renders NCT a particularly interesting antiseptic for treatment of infections with high amounts of exudate (Gottardi et al. 2014; Gottardi and Nagl 2013). Moreover, NCT kills fungi much more rapidly in the presence of organic matter than in buffer solution (Gruber et al. 2017; Lackner et al. 2015; Nagl et al. 2001).

Regarding the human defence system, the results of this study may provide evidence that single oxidants cooperate in their attack on invading microorganisms. Despite a high number of studies on highly reactive oxidants such as hypochlorite and long-lived ones such as chloramines, this aspect appears to have been underestimated up to date. Hydrogen peroxide and NCT are moderately enhanced in their bactericidal

activity if applied in combination. The relatively slowly produced singlet oxygen seems not to be responsible for this effect. Consequences are not fully foreseeable presently and may comprise the understanding of the human defence system and application of oxidants as antiseptics.

Abbreviations

ANOVA: analysis of variance; BA: bactericidal activity; Cfu: colony forming units; DMSO: dimethylsulphoxide; NCT: N-chlorotaurine; ROS: reactive oxygen species; SD: standard deviation.

Authors' contributions

JM performance of microbiological experiments, writing of the paper. MN writing of the paper, guidance of the study, statistical analysis. VX idea and concept, performance and guidance of biochemical experiments, writing of the paper. All authors read and approved the final manuscript.

Author details

¹ Division of Hygiene and Medical Microbiology, Medical University of Innsbruck, Innsbruck, Austria. ² Department of Chemistry, Faculty of Science, UNESP - São Paulo State University, Bauru, SP, Brazil. ³ Division of Hygiene and Medical Microbiology, Medical University of Innsbruck, Schöpfstr. 41, 1st Floor, 6020 Innsbruck, Austria.

Acknowledgements

We are grateful to Andrea Windisch for excellent technical assistance.

Competing interests

The authors declare that they have no competing interests.

Availability of data and materials

All relevant data are presented in the manuscript.

Funding

This study was funded by the Austrian Science Fund, Grant No. KL1459-B30 and by the Brazilian Fundação de Amparo à Pesquisa do Estado de São Paulo, Grant No. 2016/20594-5, and INCT.Bio.Nat No. 2014/50926-0.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Received: 6 April 2017 Accepted: 15 May 2017

Published online: 23 May 2017

References

- Baldry MG (1983) The bactericidal, fungicidal and sporicidal properties of hydrogen peroxide and peracetic acid. *J Appl Bacteriol* 54:417–423
- Bruch MK (2007) Toxicity and safety of topical sodium hypochlorite. *Contrib Nephrol* 154:24–38
- De Carvalho BL, Morgon NH, De Souza AR, Ximenes VF (2016) Taurine bromamine: reactivity of an endogenous and exogenous anti-inflammatory and antimicrobial amino acid derivative. *Biomolecules* 6:23
- Gottardi W, Nagl M (2002) Chemical properties of N-chlorotaurine sodium, a key compound in the human defence system. *Arch Pharm Pharm Med Chem* 335:411–421
- Gottardi W, Nagl M (2005) Chlorine covers on living bacteria: the initial step in antimicrobial action of active chlorine compounds. *J Antimicrob Chemother* 55:475–482
- Gottardi W, Nagl M (2010) N-chlorotaurine, a natural antiseptic with outstanding tolerability. *J Antimicrob Chemother* 65:399–409
- Gottardi W, Nagl M (2013) Active halogen compounds and proteinaceous material: loss of activity of topical anti-infectives by halogen consumption. *J Pharm Pharmacol* 65:213–218
- Gottardi W, Arnitz R, Nagl M (2007) N-chlorotaurine and ammonium chloride: an antiseptic preparation with strong bactericidal activity. *Int J Pharm* 335:32–40
- Gottardi W, Debabov D, Nagl M (2013) N-chloramines: a promising class of well-tolerated topical anti-infectives. *Antimicrob Agents Chemother* 57:1107–1114
- Gottardi W, Klotz S, Nagl M (2014) Superior bactericidal activity of N-bromine compounds compared to their N-chlorine analogues can be reversed under protein load. *J Appl Microbiol* 116:1427–1437
- Gottardi W, Pfeleiderer J, Nagl M (2015) The *Integral Method*, a new approach to quantify bactericidal activity. *J Microbiol Methods* 115:71–78
- Grisham MB, Jefferson MM, Melton DF, Thomas EL (1984) Chlorination of endogenous amines by isolated neutrophils. *J Biol Chem* 259:10404–10413
- Gruber M, Moser I, Nagl M, Lackner M (2017) Bactericidal and fungicidal activity of N-chlorotaurine is enhanced in cystic fibrosis sputum medium. *Antimicrob Agents Chemother* 61:e02527-16. doi:10.1128/AAC.02527-16
- Kanofsky JR, Hoogland H, Wever R, Weiss SJ (1988) Singlet oxygen production by human eosinophils. *J Biol Chem* 263:9692–9696
- Khan AU, Kasha M (1963) Luminescence of molecular oxygen in aqueous solution. *J Chem Phys* 39:2105–2106
- Kim C, Cha YN (2014) Taurine chloramine produced from taurine under inflammation provides anti-inflammatory and cytoprotective effects. *Amino Acids* 46:89–100
- Kim SO, Merchant K, Nudelman R, Beyer WF Jr, Keng T, DeAngelo J, Hausladen A, Stamler JS (2002) OxyR: a molecular code for redox-related signaling. *Cell* 109:383–396
- Kim IW, Park JM, Roh YJ, Kim JH, Choi MG, Hasan T (2016) Direct measurement of singlet oxygen by using a photomultiplier tube-based detection system. *J Photochem Photobiol, B* 159:14–23
- Klebanoff SJ (1968) Myeloperoxidase-halide-hydrogen peroxide antibacterial system. *J Bacteriol* 95:2131–2138
- Klebanoff SJ, Kettle AJ, Rosen H, Winterbourn CC, Nauseef WM (2013) Myeloperoxidase: a front-line defender against phagocytosed microorganisms. *J Leukoc Biol* 93:185–198
- Lackner M, Binder U, Reindl M, Gönül B, Fankhauser H, Mair C, Nagl M (2015) N-chlorotaurine exhibits fungicidal activity against therapy-refractory *Scedosporium species* and *Lomentospora prolificans*. *Antimicrob Agents Chemother* 59:6454–6462
- Lengfelder E, Cadenas E, Sies H (1983) Effect of DABCO (1,4-diazabicyclo[2,2,2]-octane) on singlet oxygen monomer (1270 nm) and dimol (634 and 703 nm) emission. *FEBS Lett* 164:366–370
- Lu J, Lau CW, Lee MK, Kai M (2002) Simple and convenient chemiluminescence method for the determination of melatonin. *Anal Chim Acta* 455:193–198
- Marcinkiewicz J, Kontry E (2014) Taurine and inflammatory diseases. *Amino Acids* 46:7–20
- Martini C, Hammerer-Lercher A, Zuck M, Jekle A, Debabov D, Anderson M, Nagl M (2012) Antimicrobial and anticoagulant activity of N-chlorotaurine (NCT), N, N-dichloro-2,2-dimethyltaurine (NVC-422) and N-monochloro-2,2-dimethyltaurine (NVC-612) in human blood. *Antimicrob Agents Chemother* 56:1979–1984
- Matuszak Z, Bilka MA, Reszka KJ, Chignell CF, Bilski P (2003) Interaction of singlet molecular oxygen with melatonin and related indoles. *Photochem Photobiol* 78:449–455
- Miyamoto S, Martinez GR, Medeiros MH, Di MP (2014) Singlet molecular oxygen generated by biological hydroperoxides. *J Photochem Photobiol, B* 139:24–33
- Mongkolsuk S, Helmann JD (2002) Regulation of inducible peroxide stress responses. *Mol Microbiol* 45:9–15
- Nagl M, Hengster P, Semenitz E, Gottardi W (1999) The postantibiotic effect of N-chlorotaurine on *Staphylococcus aureus*. Application in the mouse peritonitis model. *J Antimicrob Chemother* 43:805–809
- Nagl M, Lass-Flörl C, Neher A, Gunkel AR, Gottardi W (2001) Enhanced fungicidal activity of N-chlorotaurine in nasal secretion. *J Antimicrob Chemother* 47:871–874
- Nagl M, Gruber A, Fuchs A, Lell C, Lemberger EM, Borg-von Zepelin M, Würzner R (2002) Impact of N-chlorotaurine on viability and production of secreted aspartyl proteinases of *Candida* spp. *Antimicrob Agents Chemother* 46:1996–1999
- Nagl M, Nguyen VA, Gottardi W, Ulmer H, Höpfl R (2003) Tolerability and efficacy of N-chlorotaurine compared to chloramine T for treatment of chronic leg ulcers with purulent coating. *Br J Dermatol* 149:590–597
- Neher A, Nagl M, Appenroth E, Gstöttner M, Wischatta M, Reisinger F, Schindler M, Ulmer H, Stephan K (2004) Acute otitis externa: efficacy and tolerability of N-chlorotaurine, a novel endogenous antiseptic agent. *Laryngoscope* 114:850–854
- Peskin AV, Midwinter RG, Harwood DT, Winterbourn CC (2005) Chlorine transfer between glycine, taurine, and histamine: reaction rates and impact on cellular reactivity. *Free Radic Biol Med* 38:397–405
- Peskin AV, Turner R, Maghazal GJ, Winterbourn CC, Kettle AJ (2009) Oxidation of methionine to dehydromethionine by reactive halogen species generated by neutrophils. *Biochemistry* 48:10175–10182
- Sabbahi S, Alouini Z, Jemli M, Boudabbous A (2008) The role of reactive oxygen species in *Staphylococcus aureus* photoinactivation by methylene blue. *Water Sci Technol* 58:1047–1054
- Steinbeck MJ, Khan AU, Karnovsky MJ (1992) Intracellular singlet oxygen generation by phagocytosing neutrophils in response to particles coated with a chemical trap. *J Biol Chem* 267:13425–13433
- Stief TW (2003) The physiology and pharmacology of singlet oxygen. *Med Hypotheses* 60:567–572
- Stief TW, Fareed J (2000) The antithrombotic factor singlet oxygen/light (1O₂/h_ν). *Clin Appl Thromb Hemost* 6:22–30
- Stief TW, Feek U, Ramaswamy A, Kretschmer V, Renz H, Fareed J (2001) Singlet oxygen (1O₂) disrupts platelet aggregates. *Thromb Res* 104:361–370
- Teuchner B, Nagl M, Schidlbauer A, Ishiko H, Dragosits E, Ulmer H, Aoki K, Ohno S, Mizuki N, Gottardi W, Larcher C (2005) Tolerability and efficacy of N-chlorotaurine in epidemic keratoconjunctivitis—a double-blind randomized phase 2 clinical trial. *J Ocular Pharmacol Ther* 21:157–165
- Weiss SJ, Klein R, Slivka A, Wei M (1982) Chlorination of taurine by human neutrophils. *J Clin Invest* 70:598–607
- Wilkins RG, Unverdorben M (2013) Wound cleaning and wound healing: a concise review. *Adv Skin Wound Care* 26:160–163
- Winterbourn CC, Kettle AJ (2013) Redox reactions and microbial killing in the neutrophil phagosome. *Antioxid Redox Signal* 18:642–660
- Zgliczynski JM, Stelmazynska T, Domanski J, Ostrowski W (1971) Chloramines as intermediates of oxidation reaction of amino acids by myeloperoxidase. *Biochim Biophys Acta* 235:419–424